

RESVOXITROL



CLINICAL APPLICATIONS

- *Increases Antioxidant Protection*
- *Maintains Normal Inflammatory Balance*
- *Activates Nrf2, the Master Cellular Switch for Antioxidant Production*
- *Supports Cellular Longevity*
- *Enhances Mechanisms of Cellular Detoxification*

CARDIOVASCULAR HEALTH

Resvoxitol is a unique combination of phytonutrients designed to enhance antioxidant potential by stimulating nuclear factor-erythroid-2-related factor 2 (Nrf2), a dynamic pathway known to increase the production of the body's most important cellular antioxidants. This unique combination of truebroc™, turmeric, andrographis, and resveratrol provides potent free radical scavenging potential, maintaining normal inflammatory balance, and induction of intracellular antioxidant production. These powerful phytochemicals are the basis of many "superfoods" and have extensive, peer-reviewed research supporting their dramatic effect on cell health and longevity.

Overview

Nrf2 is a transcription factor that regulates the antioxidant response of the body.¹ It is the primary cellular defense against the cytotoxic effects of oxidative stress.² This key protein is found inside a cell and is inactive until it is released by an Nrf2 activator. Once activated, Nrf2 migrates into the nucleus and binds to DNA at the location of the master antioxidant regulator, the antioxidant response element (ARE).³ Studies suggest Nrf2 plays an important role in supporting the activation of cellular antioxidant systems, as well as maintaining normal inflammatory balance.^{4,5} A number of fruit and vegetable compounds have been shown to activate Nrf2 and ARE, and increase antioxidant potential.⁶⁻⁹ It is believed that these compounds have a protective effect on the body, help maintain normal inflammatory balance, and induce cellular longevity.¹⁰

Broccoli Seed Extract†

Broccoli seed extract has the highest levels of glucoraphanin (a potent Nrf2 stimulator) of any broccoli product available.¹¹ In

fact, 230 mg of broccoli seed extract is equivalent to consuming 1 ¼ cups of fresh broccoli. Glucoraphanin is a phytochemical compound that acts as the precursor to sulforaphane, one of the leading compounds in broccoli that is responsible for many of its positive benefits.^{12,13} These compounds and their mechanisms have been studied extensively and supported by more than 500 scientific publications. Glucoraphanin and sulforaphane act as direct activators of Nrf2 and potent antioxidants shown to support detoxification and maintain normal inflammatory balance.¹⁴⁻¹⁸ Studies show they work by improving hormone balance and increasing functional enzymes that support excretion of toxins from the body.¹⁹ Broccoli seed extract has been shown to remain active within the body for up to three days following consumption.²⁰

Turmeric Root Extract†

Turmeric is a bright orange-colored spice used for thousands of years in cooking and medicine. Through its antioxidant mechanisms, the active ingredient in turmeric, curcumin, helps to maintain a healthy cardiovascular system, support gastrointestinal (GI) health, and provide neuroprotective activity.²¹⁻²⁴ Curcumin has been shown to play a crucial role in maintaining normal inflammatory balance in a variety of bodily systems due to its ability to modulate Nrf2.²⁵ Studies have shown curcumin maintains normal inflammatory balance in endothelial cells and cardiac cells.²⁶⁻²⁸ Curcumin has also been found to provide antioxidant protection to the cardiovascular system by reducing oxidative stress and supporting a normal immune response.²⁹

Andrographis †

Andrographis (*Andrographis paniculata*) is an herbaceous plant native in India and Sri Lanka. It has widely been used

†These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

EFFICACY
the power of *e*

in Southeastern Asian medicine for its broad range of health-promoting properties. In recent years, andrographis has emerged as a potent activator of Nrf2.¹ Recent studies have shown that when compared to other phenolic compounds, 60 mg of andrographis induced the highest activation of Nrf2.² This activation helps reduce oxidative stress, quench free radicals, maintain endothelial health and support mitochondrial function.²⁻⁴

Resveratrol[†]

Resveratrol, is a stilbene found in many plants and red wines. It is the most well-researched stilbene. Studies of resveratrol have found it to enhance the potential of antioxidant and detoxification activity through Nrf2 activation.³⁰ Other studies have linked resveratrol to enhancing blood sugar balance, maintaining normal inflammatory balance and reducing oxidative stress.^{31, 32}

Directions

2 capsules per day or as recommended by your health care professional.

Does Not Contain

Gluten, yeast, artificial colors and flavors.

Cautions

Do not consume this product if you are pregnant or nursing. Consult your physician for further information.

Supplement Facts ^{v3}		
Serving Size 2 Capsules		
Servings Per Container 30		
2 capsules contain	Amount Per Serving	% Daily Value
Turmeric Root Extract (Standardized to contain 95% Curcuminoids)	400 mg	*
Broccoli Seed Extract (TrueBroc®) (Standardized to contain 13% Glucoraphanin)	230 mg	*
<i>trans</i> -Resveratrol (from <i>Polygonum cuspidatum</i> (Roots))	100 mg	*
<i>Andrographis paniculata</i> Extract (Leaves) (ParActin® (Bioactive 14-Neo-Andro Compound)) (Standardized to contain 50% Andrographolides)	60 mg	*
* Daily Value not established		

ID# 586060 60 Capsules

References

1. Wong SY, Tan MG, Wong PT, et al. Andrographolide induces Nrf2 and heme oxygenase 1 in astrocytes by activating p38 MAPK and ERK. *Journal of neuroinflammation* 2016;13(1):251.
2. Wu KC, McDonald PR, Liu J, et al. Screening of natural compounds as activators of the keap1-nrf2 pathway. *Planta medica* 2014;80(1):97.
3. Rajagopal S, Kumar RA, Deevi DS, et al. Andrographolide, a potential cancer therapeutic agent isolated from *Andrographis paniculata*. *Journal of Experimental therapeutics and Oncology* 2003;3(3):147-58.
4. Mishra SK, Sangwan NS, Sangwan RS. *Phcog Rev.: Plant Review Andrographis paniculata* (Kalmegh): A Review. *Pharmacognosy Reviews* 2007;1(2):283-98.
5. Li W, Khor TO, Xu C, Shen G, Jeong W-S, Yu S and Kong A-N. Activation of Nrf2-antioxidant signaling attenuates NFκB-inflammatory response and elicits apoptosis. *Biochemical pharmacology*. 2008;76:1485-1489.
6. Surh Y-J, Kundu JK and Na H-K. Nrf2 as a master redox switch in turning on the cellular signaling involved in the induction of cytoprotective genes by some chemopreventive phytochemicals. *Planta medica*. 2008;74:1526-1539.
7. Surh Y-J and Na H-K. NF-κB and Nrf2 as prime molecular targets for chemoprevention and cytoprotection with anti-inflammatory and antioxidant phytochemicals. *Genes & nutrition*. 2008;2:313-317.
8. Lee JH, Khor TO, Shu L, Su Z-Y, Fuentes F and Kong A-NT. Dietary phytochemicals and cancer prevention: Nrf2 signaling, epigenetics, and cell death mechanisms in blocking cancer initiation and progression. *Pharmacology & therapeutics*. 2013;137:153-171.
9. Gopalakrishnan A and Kong A-NT. Anticarcinogenesis by dietary phytochemicals: cytoprotection by Nrf2 in normal cells and cytotoxicity by modulation of transcription factors NF-κB and AP-1 in abnormal cancer cells. *Food and Chemical Toxicology*. 2008;46:1257-1270.

[†]These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

10. Su Z-Y, Shu L, Khor TO, Lee JH, Fuentes F and Kong A-NT. A perspective on dietary phytochemicals and cancer chemoprevention: oxidative stress, nrf2, and epigenomics. *Natural Products in Cancer Prevention and Therapy*: Springer; 2012: 133-162.
11. McWalter GK, Higgins LG, McLellan LI, Henderson CJ, Song L, Thornalley PJ, Itoh K, Yamamoto M and Hayes JD. Transcription factor Nrf2 is essential for induction of NAD (P) H: quinone oxidoreductase 1, glutathione S-transferases, and glutamate cysteine ligase by broccoli seeds and isothiocyanates. *The Journal of nutrition*. 2004;134:3499S-3506S.
12. Zhang Y, Kensler TW, Cho C-G, Posner GH and Talalay P. Anticarcinogenic activities of sulforaphane and structurally related synthetic norbornyl isothiocyanates. *Proceedings of the National Academy of Sciences*. 1994;91:3147-3150.
13. Clarke JD, Dashwood RH and Ho E. Multi-targeted prevention of cancer by sulforaphane. *Cancer letters*. 2008;269:291-304.
14. Singh AV, Xiao D, Lew KL, Dhir R and Singh SV. Sulforaphane induces caspase-mediated apoptosis in cultured PC-3 human prostate cancer cells and retards growth of PC-3 xenografts in vivo. *Carcinogenesis*. 2004;25:83-90.
15. Li Y, Zhang T, Korkaya H, Liu S, Lee H-F, Newman B, Yu Y, Clouthier SG, Schwartz SJ and Wicha MS. Sulforaphane, a dietary component of broccoli/broccoli sprouts, inhibits breast cancer stem cells. *Clinical Cancer Research*. 2010;16:2580-2590.
16. Heiss E, Herhaus C, Klimo K, Bartsch H and Gerhäuser C. Nuclear factor κ B is a molecular target for sulforaphane-mediated anti-inflammatory mechanisms. *Journal of Biological Chemistry*. 2001;276:32008-32015.
17. Thimmulappa RK, Mai KH, Srisuma S, Kensler TW, Yamamoto M and Biswal S. Identification of Nrf2-regulated genes induced by the chemopreventive agent sulforaphane by oligonucleotide microarray. *Cancer research*. 2002;62:5196-5203.
18. Morimitsu Y, Nakagawa Y, Hayashi K, Fujii H, Kumagai T, Nakamura Y, Osawa T, Horio F, Itoh K and Iida K. A sulforaphane analogue that potently activates the Nrf2-dependent detoxification pathway. *Journal of Biological Chemistry*. 2002;277:3456-3463.
19. Juge N, Mithen R and Traka M. Molecular basis for chemoprevention by sulforaphane: a comprehensive review. *Cellular and Molecular Life Sciences*. 2007;64:1105-1127.
20. Hanlon N, Coldham N, Gielbert A, Kuhnert N, Sauer MJ, King LJ and Ioannides C. Absolute bioavailability and dose-dependent pharmacokinetic behaviour of dietary doses of the chemopreventive isothiocyanate sulforaphane in rat. *British Journal of Nutrition*. 2008;99:559-564.
21. Nishinaka T, Ichijo Y, Ito M, Kimura M, Katsuyama M, Iwata K, Miura T, Terada T and Yabe-Nishimura C. Curcumin activates human glutathione S-transferase P1 expression through antioxidant response element. *Toxicology letters*. 2007;170:238-247.
22. Tapia E, Soto V, Ortiz-Vega KM, Zarco-Márquez G, Molina-Jijón E, Cristóbal-García M, Santamaría J, García-Niño WR, Correa F and Zazueta C. Curcumin induces Nrf2 nuclear translocation and prevents glomerular hypertension, hyperfiltration, oxidant stress, and the decrease in antioxidant enzymes in 5/6 nephrectomized rats. *Oxidative medicine and cellular longevity*. 2012;2012.
23. Aggarwal BB and Harikumar KB. Potential therapeutic effects of curcumin, the anti-inflammatory agent, against neurodegenerative, cardiovascular, pulmonary, metabolic, autoimmune and neoplastic diseases. *The international journal of biochemistry & cell biology*. 2009;41:40-59.
24. Zhang F, Altorki NK, Mestre JR, Subbaramaiah K and Dannenberg AJ. Curcumin inhibits cyclooxygenase-2 transcription in bile acid-and phorbol ester-treated human gastrointestinal epithelial cells. *Carcinogenesis*. 1999;20:445-451.
25. Balogun E, Hoque M, Pengfei G, Killeen E, Green CJ, Foresti R, Jawed A and Motterlini R. Curcumin activates the haem oxygenase-1 gene via regulation of Nrf2 and the antioxidant-responsive element. *Biochemical Journal*. 2003;371:887-895.
26. Motterlini R, Foresti R, Bassi R and Green CJ. Curcumin, an antioxidant and anti-inflammatory agent, induces heme oxygenase-1 and protects endothelial cells against oxidative stress. *Free Radical Biology and Medicine*. 2000;28:1303-1312.

27. Fiorillo C, Becatti M, Pensalfini A, Cecchi C, Lanzilao L, Donzelli G, Nassi N, Giannini L, Borch E and Nassi P. Curcumin protects cardiac cells against ischemia-reperfusion injury: effects on oxidative stress, NF- κ B, and JNK pathways. *Free Radical Biology and Medicine*. 2008;45:839-846.
28. Yeh C-H, Chen T-P, Wu Y-C, Lin Y-M and Lin PJ. Inhibition of NF κ B Activation with Curcumin Attenuates Plasma Inflammatory Cytokines Surge and Cardiomyocytic Apoptosis Following Cardiac Ischemia/Reperfusion 1. *Journal of Surgical Research*. 2005;125:109-116.
29. Diaz MN, Frei B, Vita JA and Keaney Jr JF. Antioxidants and atherosclerotic heart disease. *New England Journal of Medicine*. 1997;337:408-416.
30. Kode A, Rajendrasozhan S, Caito S, Yang S-R, Megson IL and Rahman I. Resveratrol induces glutathione synthesis by activation of Nrf2 and protects against cigarette smoke-mediated oxidative stress in human lung epithelial cells. *American Journal of Physiology-Lung Cellular and Molecular Physiology*. 2008;294:L478-L488.
31. Su H-C, Hung L-M and Chen J-K. Resveratrol, a red wine antioxidant, possesses an insulin-like effect in streptozotocin-induced diabetic rats. *American Journal of Physiology-Endocrinology and Metabolism*. 2006;290:E1339-E1346.
32. Martí AR, Villegas I, La Casa C and de la Lastra CA. Resveratrol, a polyphenol found in grapes, suppresses oxidative damage and stimulates apoptosis during early colonic inflammation in rats. *Biochemical pharmacology*. 2004;67:1399-1410.