

# LIQUID Vitamin D3 with K2



## CLINICAL APPLICATIONS

- Supports Healthy Blood Circulation
- Promotes Bone Health and Proper Calcium Storage
- Maintains Healthy Cardiometabolic Function and Supports Blood Sugar Balance Already Within Normal Levels
- Boosts Immune Function

## ESSENTIAL VITAMINS

Well-established research has focused on the synergistic relationship between vitamin K2 and vitamin D3 for bone and cardiovascular health.<sup>1</sup> A group of naturally occurring and structurally similar fat-soluble vitamins, vitamin K is essential for the proper utilization of calcium. Through its activation of the protein osteocalcin, vitamin K helps to bind newly absorbed calcium to the mineral matrix in bone. In addition, vitamin K has been found to help maintain bone mineral density by decreasing the activity of osteoclasts, a cell that breaks down the bone matrix.<sup>2</sup> Vitamin K also provides critical cardiovascular protection by helping to activate matrix Gla protein (MGP), an inhibitor of circulatory calcification.<sup>3,4</sup> Thus, vitamin K and vitamin D not only share similar qualities but they also act synergistically with one another within the body.<sup>5</sup> Liquid Vitamin D3 with K2 utilizes a medium-chain triglyceride (MCT) delivery system, providing 10 mcg of MenaQ7<sup>®</sup> PRO, the most widely studied form of vitamin K2 as MK-7, and 1,000 IU of vitamin D3 per serving. Natural medium-chain triglycerides have unique physical properties that allow them to be readily absorbed without additional emulsifying agents. MCTs are a great way to deliver pure, tasteless and odorless liquid vitamin K2 and D3 that combine easily with any beverage.

### Overview

While vitamin D has long been known to assist calcium absorption, it is vitamin K, through its carboxylation of osteocalcin, which guides this calcium to bones and prevents their absorption into organs, joint spaces and arteries.<sup>6</sup> Vitamin K occurs in two main forms: K1 (phylloquinone), derived from

foods such as green leafy vegetables and K2 (menaquinone), which is a group of related compounds differentiated by their side chains. Numerous studies have shown that vitamin K2 is the more bioavailable form of the nutrient and more powerfully influences bone building than K1.<sup>7</sup> In addition, though both reach the liver, most of the K1 is used for purposes of coagulation, with little left over to support the body's needs elsewhere.<sup>8</sup> The profoundly different degrees of bioavailability between K1 and K2 is due to their differences in structure; only 10-20% of vitamin K1 that is absorbed from food even reaches the circulation, while the long side chain of vitamin K2 allows it to bind with fat particles in circulation and facilitate its arrival at soft tissue, bones and arteries. There are two forms of vitamin K2 commonly used in supplements: MK-4 and MK-7. The MK-7 form has been shown to have six times the activity of MK-4 in the blood. MK-7 has also been found to remain in the blood approximately nine times as long as MK-4 (eight hours versus 72 hours), making it the optimal form of K2.<sup>9</sup>

The addition of MenaQ7<sup>®</sup> PRO to this formula is backed by extensive research. Dr. Leon Schurgers, world-renowned expert in vitamin K2 as MK-7 research for cardiovascular and bone health, and his team of research scientists have conducted over 15 clinical trials on MenaQ7<sup>®</sup> PRO through the University of Maastricht.

### Vitamin K Depletion<sup>†</sup>

Although most people consume adequate dietary vitamin K to maintain sufficient blood clotting, most do *not* consume enough to meet cardiovascular and bone health needs. In fact, approximately 70% of the Western population is deficient in

<sup>†</sup> These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

vitamin K2. Compromised intestinal absorption can also lead to insufficient K2 levels, leaving calcium available to be exported out of bone and into other tissues. Other medications, such as antibiotics, cholesterol-lowering medications and laxatives, have also been found to contribute to a deficiency of vitamin K.<sup>10</sup>

### Bone Health<sup>†</sup>

Building and maintaining healthy bones requires a number of key nutrients, including vitamin K, for the proper binding of calcium to the bone matrix. A 2005 study from northern Finland found that those with greater levels of vitamin K-carboxylated osteocalcin had stronger bones than those with lower levels of the protein. Vitamin K2 alone does not improve bone mineral content, it also requires vitamin D. The synthesis of osteocalcin and dietary absorption of calcium are vitamin D dependent processes.<sup>11</sup> A Japanese study found superior bone health among women who were frequent MK-7-rich natto eaters than those who were not.<sup>12</sup> Another randomized study which split 172 women into a vitamin K2 group, a vitamin D3 group, a vitamin K2 and D3 group, and a placebo group for two years found that the combination of vitamin D3 and K2 had the most benefits for supporting bone health among the groups.<sup>13</sup>

### Cardiovascular Health and Blood Sugar Balance<sup>†</sup>

Vitamin K plays a key role in supporting the cardiovascular system and blood sugar balance already within normal levels. In a large population study, researchers found that those who consumed high amounts of K2 had significantly better cardiovascular health markers compared to those given vitamin K1.<sup>14</sup> Studies have also shown vitamin K supports healthy blood sugar metabolism.<sup>15,16</sup>

### Additional Health Benefits<sup>†</sup>

Vitamin K2 has additional health benefits beyond bone and cardiometabolic health benefits. Vitamin K2 has been shown to support mitochondrial function, muscle and nervous tissue.<sup>17,18</sup> In a 12 week study of 100 people with nerve pain, 100mcg of Vitamin K2 MK-7 had significant improvement in four variables of assessed pain.<sup>18</sup> Both vitamin D and K impart immune-modulating effects. Vitamins D and K are associated with stronger immune function, balanced inflammatory response and improved longevity.<sup>11,19,20,21</sup>

### Directions

1 drop (0.025 mL) per day or as recommended by your health care professional.

### Does Not Contain

Gluten, corn, yeast, artificial colors or flavors.

<b>Supplement Facts</b> <sup>V3</sup>		
Serving Size 1 Drop (0.025 mL)		
Servings Per Container About 1200		
	Amount Per Serving	% Daily Value
Vitamin D (D3 as Cholecalciferol)	25 mcg (1,000 IU)	125%
Vitamin K2 (as Menaquinone-7 (MK-7)) (MenaQ7 <sup>®</sup> PRO)	10 mcg	*

\* Daily Value not established.

Other Ingredients: Medium Chain Triglycerides.

**ID# 870001 1 fl oz (30 ml)**



MenaQ7<sup>®</sup> PRO is a registered trademark of NattoPharma, Norway.

### Cautions

If you are pregnant or nursing, consult your physician before taking this product. All forms of vitamin K may interact with blood thinning medications. If you are taking such medicines, please consult your physician before taking this product.

<sup>†</sup> These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

## References

1. Schurgers LJ, Spronk HM, Soute BA, Schiffrers PM, DeMey JG, Vermeer C. Regression of warfarin-induced medial elastocalcinosis by high intake of vitamin K in rats. *Blood*. 2007;109(7):2823-31.
2. Weber P. Management of osteoporosis: is there a role for vitamin K? *Int J Vitam Nutr Res*. 1997;67(5):350-6.
3. Geleijnse JM, et al. Dietary intake of menaquinone is associated with a reduced risk of coronary heart disease: the Rotterdam Study. *J Nutr*. 2004;134(11):3100-5.
4. Beulens JW, High dietary menaquinone intake is associated with reduced coronary calcification. *Atherosclerosis*. 2009;203(2):489-93. Epub 2008 Jul 19.
5. Kidd PM. Vitamins D and K as pleiotropic nutrients: clinical importance to the skeletal and cardiovascular systems and preliminary evidence for synergy. *Altern Med Rev*. 2010;15(3):199-222.
6. Plaza SM, Lamson DW. Vitamin K2 in bone metabolism and osteoporosis. *Altern Med Rev*. 2005;10(1):24-35.
7. Schurgers LJ, Teunissen KJ, Hamulyák K, Knapen MH, Vik H, Vermeer C. Vitamin K-containing dietary supplements: comparison of synthetic vitamin K1 and natto-derived menaquinone. *Blood*. 2007;109(8): 3279-83. Epub 2006 Dec 7.
8. Shearer MJ. Vitamin K metabolism and nutriture. *Blood Rev*. 1992;6(2):92-104.
9. Unpublished clinical studies, NattoPharma. On file
10. <http://umm.edu/health/medical/altmed/supplement-depletion-links/drugs-that-deplete-vitamin-k>
11. Marles RJ, Roe AL, & Oketch-Rabah HA. (2017). US Pharmacopeial Convention safety evaluation of menaquinone-7, a form of vitamin K. *Nutrition Reviews*, 75(7), 553–578. <https://doi.org/10.1093/nutrit/nux022>.
12. Kaneki M, Hodges SJ, Hosoi T, et al. Japanese fermented soybean food as the major determinant of the large geographic difference in circulating levels of vitamin K2: possible implications for hip-fracture risk. *Nutrition*. 2001; 17(4):315-21.
13. Ushiroyama T, Ikeda A, Ueki M. Effect of continuous combined therapy with vitamin K(2) and vitamin D(3) on bone mineral density and coagulofibrinolysis function in postmenopausal women. *Maturitas*. 2002; 41(3):211-21.
14. Geleijnse JM, Vermeer C, Grobbee DE, et al. Dietary intake of menaquinone is associated with a reduced risk of coronary heart disease: the Rotterdam Study. *J Nutr*. 2004; 134(11):3100-5.
15. Beulens JW, van der A DL, Grobbee DE, Sluijs I, Spijkerman AM, van der Schouw YT. Dietary phylloquinone and menaquinones intakes and risk of type 2 diabetes. *Diabetes Care*. 2010; 33(8):1699-705.
16. Choi HJ, Yu J, Choi H, et al. Vitamin K2 supplementation improves insulin sensitivity via osteocalcin metabolism: a placebo-controlled trial. *Diabetes Care*. 2011; 34(9):e147.
17. Koziół-Kozakowska A, & Maresz K. (2022). The Impact of Vitamin K2 (Menaquinones) in Children's Health and Diseases: A Review of the Literature. *Children*, 9(1), 78. <https://doi.org/10.3390/children9010078>.
18. Mehta DS, Dound YA, Jadhav SS, Bhave AA, Milind Devale & Vaidya ADB. (2018). A Novel Potential Role of Vitamin K2-7 in Relieving Peripheral Neuropathy. *Journal of Pharmacology and Pharmacotherapeutics* (Print), 9(4), 180–185. [https://doi.org/10.4103/jpp.jpp\\_72\\_18](https://doi.org/10.4103/jpp.jpp_72_18).
19. Beulens JWJ, van der A DL, Grobbee DE, Sluijs I, Spijkerman AMW, & van der Schouw YT. (2010). Dietary Phylloquinone and Menaquinones Intakes and Risk of Type 2 Diabetes. *Diabetes Care*, 33(8), 1699–1705. <https://doi.org/10.2337/dc09-2302>.
20. Heath A, Kim I, Hodge A, English D, & Muller D. (2019). Vitamin D Status and Mortality: A Systematic Review of Observational Studies. *International Journal of Environmental Research and Public Health*, 16(3), 383. <https://doi.org/10.3390/ijerph16030383>.
21. Iijima H, Shinzaki S, Takehara T. The importance of vitamins D and K for the bone health and immune function in inflammatory bowel disease. *Curr Opin Clin Nutr Metab Care*. 2012; 15(6):635-40.